

T6 SPINAL SOLITARY BREAST METASTASIS



UPMC
HEALTH
SYSTEM

University of Pittsburgh Medical Center CyberKnife® Team:

Surgeon: Peter C. Gerszten, M.D., M.P.H.

Radiation Oncologist: Steven A. Burton, M.D.

Medical Physicist: Cihat Ozhasoglu, Ph.D.

Radiation Therapist: William J. Vogel, R.T.(T.)

Nurse: Annette E. Quinn, R.N.

CyberKnife Center: Shadyside Hospital
University of Pittsburgh
Medical Center
Pittsburgh, PA

T6 SPINAL SOLITARY BREAST METASTASIS

DEMOGRAPHICS

Sex: F
Age: 56
Histology: Breast Metastasis to T6

CLINICAL HISTORY


Referred by: Medical Oncology
Previous Treatment: External beam radiation of 30 Gy in 10 fractions to the T6 vertebral body

Case History

The patient originally was diagnosed with a T2, N0 infiltrating ductal carcinoma of the right breast. Her cancer was treated with segmental resection. This was followed by radiation therapy to the breast to 61.2 Gy and four cycles of adjuvant 5-FU and methotrexate. Six years later, she presented to her medical oncologist with complaints of back pain. MR imaging revealed a solitary destructive lesion of the T6 vertebral body. Further workup failed to demonstrate other areas of metastatic disease. This solitary metastatic lesion was treated with external beam radiation in ten fractions to 30 Gy with temporary improvement in her symptoms.

However, a month later, persistent symptoms of pain prevented the patient from returning to work and interfered with her activities of daily living. She was therefore referred to the CyberKnife® Spine Center for further evaluation.


Her MRI showed tumor progression with significant compression of the spinal cord at the T6 level. She had no neurological deficits.



Pretreatment MR: Sagittal T2 weighted and axial T1 weighted gadolinium-enhanced MRI reveals a pathologic compression fracture with significant spinal canal compromise.

CyberKnife Treatment Rationale

The treatment of both malignant as well as benign tumors of the spine using CyberKnife radiosurgery began in 1997. Treatment of spine lesions using single fraction radiosurgery has been a successful treatment strategy at UPMC over the past four years.^{1,2,3}



Pretreatment CT showing the outlined tumor at the T6 level. This image set was used for treatment planning and stereotactic radiosurgical targeting on the CyberKnife System.

TREATMENT DETAILS

Tumor Volume: 10.3 cc
Imaging Technique(s): CT, T2-MRI, MRI+C
Rx Dose & Isodose: 16 Gy to 80%
Conformality Index: 1.14
Tumor Coverage: 72.7% of PTV
Number of Beams: 103



Fractions / Treatment Time: 1 fraction in 65 minutes
Path Template: 1 path 900_1000 mm
Tracking Method: Tracking with 5 fiducials
Collimator(s): 15 mm

Planning Process and Goals


The tumor target volume of 10.3 cc was contoured to include the tumor at the level of the T6 vertebral body. Surrounding critical structures, including the spinal cord and esophagus, were contoured to minimize dose to those radiosensitive structures. The 80% isodose line represents the prescribed dose of 16 Gy to the tumor. The prescription isodose line covered 72.7% of the planning target volume with a conformality index of 1.14.

Treatment Delivery


The treatment utilized 103 separately targeted beams with the 15 mm collimator. The tumor volume measured 10.3 cc and only 0.3 cc of the spinal cord received greater than 8 Gy, consistent with the low conformality index. The patient reported no adverse side effects from the treatment delivery.



Axial and coronal planning images with the tumor, isodose curves and critical structures outlined. Note the highly conformal dose distribution avoids the critical spinal cord and esophagus at the level of T6 as seen in the axial view.



Anterior-posterior rendering of the CyberKnife® System's beam positions for this treatment.



Dose Volume Histogram (DVH) for tumor and critical structures - cord and esophagus.